


Reactions of Benzene & Its Derivatives


Reactions of Benzene

The most characteristic reaction of aromatic compounds is substitution at a ring carbon:

Halogenation:




Nitration:




Reactions of Benzene

ORGANIC LECTURE SERIES


Sulfonation:




Alkylation:



Acylation:




3



Electrophilic Aromatic Substitution

- **Electrophilic aromatic substitution:** a reaction in which a hydrogen atom of an aromatic ring is replaced by an electrophile




- In this section:
 - several common types of electrophiles
 - how each is generated
 - the mechanism by which each replaces hydrogen

5

EAS: General Mechanism



- A general mechanism



- Key question: What is the electrophile and how is it generated?


6

Electrophilic Aromatic Substitution (EAS)




Chlorination

Step 1: formation of a chloronium ion




Step 2: attack of the chloronium ion on the ring




Chlorination

Step 3: proton transfer regenerates the aromatic character of the ring



9


Bromination



This is the general method for
Substitution of halogen onto a benzene ring
(CANNOT be halogenated by Free Radical Mechanism)


10

Bromination-Why not addn of Br₂?




Nitration

- Generation of the nitronium ion, NO₂⁺
 - Step 1: proton transfer to nitric acid




- Step 2: loss of H₂O gives the nitronium ion, a very strong electrophile




Nitration

ORGANIC LECTURE SERIES

Step 1: attack of the nitronium ion (an electrophile) on the aromatic ring (a nucleophile)



Step 2: proton transfer regenerates the aromatic ring




13

Nitration

ORGANIC LECTURE SERIES


- A particular value of nitration is that the nitro group can be reduced to a 1° amino group



14

Sulfonation

- Carried out using concentrated sulfuric acid containing dissolved sulfur trioxide




(SO_3 in H_2SO_4 is sometimes called "fuming" sulfuric acid.)

15

Friedel-Crafts Alkylation

- Friedel-Crafts alkylation forms a new C-C bond between an aromatic ring and an alkyl group




The electrophilic partner is a carbocation;
it will arrange to the most stable ion: allylic > 3° > 2° > 1°

16


Friedel-Crafts Alkylation

ORGANIC LECTURE SERIES


Step 1: formation of an alkyl cation as an ion pair



Step 2: attack of the alkyl cation on the aromatic ring



Step 3: proton transfer regenerates the aromatic ring




17

Friedel-Crafts Alkylation

ORGANIC LECTURE SERIES

There are two major limitations on Friedel-Crafts alkylations:


1. carbocation rearrangements are common:




18

Friedel-Crafts Alkylation


2. F-C alkylation fails on benzene rings bearing one or more of these strongly **electron-withdrawing groups**



When Y Equals Any of These Groups, the Benzene Ring Does Not Undergo Friedel-Crafts Alkylation




19



20

The "De-activation" of Aromatic Systems




If the FG is an e⁻ withdrawing substituent, then the ring system becomes more electron poor and is said to be "deactivated" towards electrophilic aromatic substitution. EAS occurs at a slower rate.

Note: deactivation refers to the **rate** of EAS

21

Friedel-Crafts Acylation

- Friedel-Crafts **acylation** forms a new C-C bond between a benzene ring and an acyl group:




22

Friedel-Crafts Acylation

ORGANIC LECTURE SERIES

- The electrophile is an **acylium ion**




23

Friedel-Crafts Acylation

ORGANIC LECTURE SERIES

- an acylium ion is a resonance hybrid of two major contributing structures




- F-C acylations are free of a major limitation of F-C alkylations; **acylium ions do not rearrange.**

24

Friedel-Crafts Acylation

ORGANIC LECTURE SERIES


A special value of F-C acylations is preparation of **unrearranged** alkylbenzenes:



25

Di- and Polysubstitution

ORGANIC LECTURE SERIES



26

Di- and Polysubstitution

Orientation on nitration of monosubstituted benzenes:

Substituent	ortho	meta	para	ortho + para	meta
—OCH ₃	44	-	55	99	trace
—CH ₃	58	4	38	96	4
—Cl	70	-	30	100	trace
—Br	37	1	62	99	1
—COOH	18	80	2	20	80
—CN	19	80	1	20	80
—NO ₂	6.4	93.2	0.3	6.7	93.2

27

Di- and Polysubstitution

- **Orientation:**


- certain substituents direct preferentially to ortho & para positions; others to meta positions
- substituents are classified as either **ortho-para directing** or **meta directing** toward further substitution

28


• Rate

- certain substituents cause the rate of a second substitution to be greater than that for benzene itself; others cause the rate to be lower
- substituents are classified as **activating** or **deactivating** toward further substitution

29



If the FG is an e⁻ donating substituent, then the ring system becomes more electron rich and is said to be "activated" towards electrophilic aromatic substitution. EAS is at a faster rate.




If the FG is an e⁻ withdrawing substituent, then the ring system becomes more electron poor and is said to be "deactivated" towards electrophilic aromatic substitution. EAS occurs at a slower rate.

30


Di- and Polysubstitution

ORGANIC LECTURE SERIES

– **-OCH₃ is ortho-para directing:**



– **-CO₂H is meta directing**



31

Di- and Polysubstitution


ORGANIC LECTURE SERIES

Ortho-para Directing	Strongly activating	$-\ddot{\text{N}}\text{H}_2$	$-\ddot{\text{N}}\text{HR}$	$-\ddot{\text{N}}\text{R}_2$	$-\ddot{\text{O}}\text{H}$	$-\ddot{\text{O}}\text{R}$
	Moderately activating	$-\ddot{\text{N}}\text{H}\overset{\text{O}}{\parallel}\text{CR}$	$-\ddot{\text{N}}\text{H}\overset{\text{O}}{\parallel}\text{CAr}$	$-\ddot{\text{O}}\overset{\text{O}}{\parallel}\text{CR}$	$-\ddot{\text{O}}\overset{\text{O}}{\parallel}\text{CAr}$	
	Weakly activating	$-\text{R}$				
	Weakly deactivating	$-\ddot{\text{F}}:$	$-\ddot{\text{Cl}}:$	$-\ddot{\text{Br}}:$	$-\ddot{\text{I}}:$	
Meta Directing	Moderately deactivating	$-\overset{\text{O}}{\parallel}\text{CH}$	$-\overset{\text{O}}{\parallel}\text{CR}$	$-\overset{\text{O}}{\parallel}\text{COH}$	$-\overset{\text{O}}{\parallel}\text{COR}$	
		$-\overset{\text{O}}{\parallel}\text{CNH}_2$	$-\text{SO}_3\text{H}$	$-\text{C}\equiv\text{N}$		
	Strongly deactivating	$-\text{NO}_2$	$-\text{NH}_3^+$	$-\text{CF}_3$	$-\text{CCl}_3$	

32

Di- and Polysubstitution ORGANIC LECTURE SERIES

the order of steps is important:



33

Theory of Directing Effects ORGANIC LECTURE SERIES

- The rate of EAS is limited by the slowest step in the reaction
- For almost every EAS, **the rate-determining step is attack of E⁺** on the aromatic ring to give a resonance-stabilized cation intermediate
- The **more stable this cation intermediate, the faster the rate-determining step** and the faster the overall reaction

34

Theory of Directing Effects


- For ortho-para directors, ortho-para attack forms a more stable cation than meta attack
 - ortho-para products are formed faster than meta products
- For meta directors, meta attack forms a more stable cation than ortho-para attack
 - meta products are formed faster than ortho-para products

35

Theory of Directing Effects

Nitration of anisole


$-\text{OCH}_3$; examine the meta attack:



36

Nitration of anisole

$-\text{OCH}_3$: examine the ortho-para attack:




This resonance structure accounts for the selectivity

37

Theory of Directing Effects

Nitration of benzoic acid

$-\text{NO}_2$; examine the meta attack:




38

Nitration of benzoic acid

ORGANIC LECTURE SERIES

$-\text{NO}_2$: assume ortho-para attack:



This resonance structure accounts for the selectivity

39

Activating-Deactivating

ORGANIC LECTURE SERIES

- Any resonance effect, such as that of $-\text{NH}_2$, $-\text{OH}$, and $-\text{OR}$, that delocalizes the positive charge on the cation intermediate lowers the activation energy for its formation, and has an **activating** effect toward further EAS
- Any resonance effect, such as that of $-\text{NO}_2$, $-\text{CN}$, $-\text{CO}$, and $-\text{SO}_3\text{H}$, that decreases electron density on the ring **deactivates** the ring toward further EAS

40

Activating-Deactivating

- Any inductive effect, such as that of -CH₃ or other alkyl group, that releases electron density toward the ring **activates** the ring toward further EAS
- Any inductive effect, such as that of halogen, -NR₃⁺, -CCl₃, or -CF₃, that **decreases electron density** on the ring **deactivates** the ring toward further EAS

41


Di- and Polysubstitution

- **Generalizations:**
 - alkyl, phenyl, and all other substituents in which the atom bonded to the ring has an unshared pair of electrons are **ortho-para directing**; all other substituents are meta directing
 - all ortho-para directing groups except the halogens are **activating** toward further substitution;
 - the halogens are **weakly deactivating**

42


Activating-Deactivating

- for the **halogens**, the inductive and resonance effects run counter to each other, but the former is somewhat stronger
- the net effect is that halogens are deactivating but ortho-para directing




43

Di- and Polysubstitution

Ortho-para Directing	Strongly activating	$-\ddot{\text{N}}\text{H}_2$	$-\ddot{\text{N}}\text{HR}$	$-\ddot{\text{N}}\text{R}_2$	$-\ddot{\text{O}}\text{H}$	$-\ddot{\text{O}}\text{R}$
	Moderately activating	$-\ddot{\text{N}}\text{H}\overset{\text{O}}{\parallel}\text{CR}$	$-\ddot{\text{N}}\text{H}\overset{\text{O}}{\parallel}\text{CAr}$	$-\ddot{\text{O}}\overset{\text{O}}{\parallel}\text{CR}$	$-\ddot{\text{O}}\overset{\text{O}}{\parallel}\text{CAr}$	
	Weakly activating	$-\text{R}$				
	Weakly deactivating	$-\ddot{\text{F}}:$	$-\ddot{\text{Cl}}:$	$-\ddot{\text{Br}}:$	$-\ddot{\text{I}}:$	
Meta Directing	Moderately deactivating	$-\overset{\text{O}}{\parallel}\text{CH}$	$-\overset{\text{O}}{\parallel}\text{CR}$	$-\overset{\text{O}}{\parallel}\text{COH}$	$-\overset{\text{O}}{\parallel}\text{COR}$	
		$-\overset{\text{O}}{\parallel}\text{CNH}_2$	$-\text{SO}_3\text{H}$	$-\text{C}\equiv\text{N}$		
	Strongly deactivating	$-\text{NO}_2$	$-\text{NH}_3^+$	$-\text{CF}_3$	$-\text{CCl}_3$	

44


Benzodiazepins



- 1) Sedative-hypnotic
- 2) Anticonvulsant
- 3) Muscle relaxant
- 4) Anxiolytic


45

Retrosynthetic Analysis



46


Short Problem Using EAS: the synthesis of p-Aminochlorobenzene



Separate o from p


47

The Synthesis of the amide section:




48

Friedel Crafts Acylation:





Amide is activating & o p directing

49




50

Xanax (Alprazolam)



Haloperidol


treatment of schizophrenia and acute psychotic states and delirium.



Chlorpromazine (Thorazine)


The introduction (1950) of chlorpromazine into clinical use has been described as the single greatest advance in psychiatric care, dramatically improving the prognosis of patients in psychiatric hospitals worldwide the availability of antipsychotic drugs curtailed indiscriminate use of electroconvulsive therapy and psychosurgery, and was one of the driving forces behind the deinstitutionalization movement.

Medicinal Chemistry




53

Medicinal Chemistry




54

Michael Reaction in Context




55

Dieckmann Condensation in Context




56

Medicinal Chemistry



57

Medicinal Chemistry



58

